

University of Groningen

Revealing the genetic roots of obesity and type 2 diabetes

Vliet-Ostaptchouk, Jana Vladimirovna van

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2010

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Vliet-Ostaptchouk, J. V. V. (2010). *Revealing the genetic roots of obesity and type 2 diabetes*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

References

1. McCarthy MI, Abecasis GR, Cardon LR, Goldstein DB, Little J, Ioannidis JP, et al. Genome-wide association studies for complex traits: consensus, uncertainty and challenges. *Nat Rev Genet.* 2008 May;9(5):356-69.
2. Doria A, Patti ME, Kahn CR. The emerging genetic architecture of type 2 diabetes. *Cell Metab.* 2008 Sep;8(3):186-200.
3. Loos RJ. Recent progress in the genetics of common obesity. *Br J Clin Pharmacol.* 2009 Dec;68(6):811-29.
4. McCarthy MI, Zeggini E. Genome-wide association studies in type 2 diabetes. *Curr Diab Rep.* 2009 Apr;9(2):164-71.
5. Walley AJ, Asher JE, Froguel P. The genetic contribution to non-syndromic human obesity. *Nat Rev Genet.* 2009 Jul;10(7):431-42.
6. Farooqi IS, O'Rahilly S. Monogenic obesity in humans. *Annu Rev Med.* 2005;56:443-58.
7. Chambers JC, Elliott P, Zabaneh D, Zhang W, Li Y, Froguel P, et al. Common genetic variation near MC4R is associated with waist circumference and insulin resistance. *Nat Genet.* 2008 Jun;40(6):716-8.
8. Loos RJ, Lindgren CM, Li S, Wheeler E, Zhao JH, Prokopenko I, et al. Common variants near MC4R are associated with fat mass, weight and risk of obesity. *Nat Genet.* 2008 Jun;40(6):768-75.
9. Vaxillaire M, Froguel P. Monogenic diabetes in the young, pharmacogenetics and relevance to multifactorial forms of type 2 diabetes. *Endocr Rev.* 2008 May;29(3):254-64.
10. Frayling TM. Genome-wide association studies provide new insights into type 2 diabetes aetiology. *Nat Rev Genet.* 2007 Sep;8(9):657-62.
11. Legry V, Cotel D, Ferrieres J, Arveiler D, Andrieux N, Bingham A, et al. Effect of an FTO polymorphism on fat mass, obesity, and type 2 diabetes mellitus in the French MONICA Study. *Metabolism.* 2009 Jul;58(7):971-5.
12. van Vliet-Ostaptchouk JV, Onland-Moret NC, van Haeften TW, Franke L, Elbers CC, Shiri-Sverdlov R, et al. HHEX gene polymorphisms are associated with type 2 diabetes in the Dutch Breda cohort. *Eur J Hum Genet.* 2008 May;16(5):652-6.
13. van Vliet-Ostaptchouk JV, Shiri-Sverdlov R, Zhernakova A, Strengman E, van Haeften TW, Hofker MH, et al. Association of variants of transcription factor 7-like 2 (TCF7L2) with susceptibility to type 2 diabetes in the Dutch Breda cohort. *Diabetologia.* 2007 Jan;50(1):59-62.
14. Zeggini E, Scott LJ, Saxena R, Voight BF, Marchini JL, Hu T, et al. Meta-analysis of genome-wide association data and large-scale replication identifies additional susceptibility loci for type 2 diabetes. *Nat Genet.* 2008 May;40(5):638-45.
15. Saxena R, Voight BF, Lyssenko V, Burt NP, de Bakker PI, Chen H, et al. Genome-wide association analysis identifies loci for type 2 diabetes and triglyceride levels. *Science.* 2007 Jun 1;316(5829):1331-6.
16. Scott LJ, Mohlke KL, Bonnycastle LL, Willer CJ, Li Y, Duren WL, et al. A genome-wide association study of type 2 diabetes in Finns detects multiple susceptibility variants. *Science.* 2007 Jun 1;316(5829):1341-5.
17. Zeggini E, Weedon MN, Lindgren CM, Frayling TM, Elliott KS, Lango H, et al. Replication of genome-wide association signals in UK samples reveals risk loci for type 2 diabetes. *Science.* 2007 Jun 1;316(5829):1336-41.
18. Helgadottir A, Thorleifsson G, Magnusson KP, Gretarsdottir S, Steinthorsdottir V, Manolescu A, et al. The same sequence variant on 9p21 associates with myocardial infarction, abdominal aortic aneurysm and intracranial aneurysm. *Nat Genet.* 2008 Feb;40(2):217-24.
19. McPherson R, Pertsemlidis A, Kavaslar N, Stewart A, Roberts R, Cox DR, et al. A common allele on chromosome 9 associated with coronary heart disease. *Science.* 2007 Jun 8;316(5830):1488-91.

20. Silander K, Tang H, Myles S, Jakkula E, Timpson NJ, Cavalli-Sforza L, et al. Worldwide patterns of haplotype diversity at 9p21.3, a locus associated with type 2 diabetes and coronary heart disease. *Genome Med.* 2009;1(5):51.
21. Broadbent HM, Peden JF, Lorkowski S, Goel A, Ongen H, Green F, et al. Susceptibility to coronary artery disease and diabetes is encoded by distinct, tightly linked SNPs in the ANRIL locus on chromosome 9p. *Hum Mol Genet.* 2008 Mar 15;17(6):806-14.
22. Manolio TA, Collins FS, Cox NJ, Goldstein DB, Hindorff LA, Hunter DJ, et al. Finding the missing heritability of complex diseases. *Nature.* 2009 Oct 8;461(7265):747-53.
23. Ioannidis JP, Thomas G, Daly MJ. Validating, augmenting and refining genome-wide association signals. *Nat Rev Genet.* 2009 May;10(5):318-29.
24. Reich DE, Lander ES. On the allelic spectrum of human disease. *Trends Genet.* 2001 Sep;17(9):502-10.
25. Redon R, Ishikawa S, Fitch KR, Feuk L, Perry GH, Andrews TD, et al. Global variation in copy number in the human genome. *Nature.* 2006 Nov 23;444(7118):444-54.
26. Tuzun E, Sharp AJ, Bailey JA, Kaul R, Morrison VA, Pertz LM, et al. Fine-scale structural variation of the human genome. *Nat Genet.* 2005 Jul;37(7):727-32.
27. Bochukova EG, Huang N, Keogh J, Henning E, Purmann C, Blaszczyk K, et al. Large, rare chromosomal deletions associated with severe early-onset obesity. *Nature.* 2009 Dec 6.
28. Thorleifsson G, Walters GB, Gudbjartsson DF, Steinthorsdottir V, Sulem P, Helgadóttir A, et al. Genome-wide association yields new sequence variants at seven loci that associate with measures of obesity. *Nat Genet.* 2009 Jan;41(1):18-24.
29. Willer CJ, Speliotes EK, Loos RJ, Li S, Lindgren CM, Heid IM, et al. Six new loci associated with body mass index highlight a neuronal influence on body weight regulation. *Nat Genet.* 2009 Jan;41(1):25-34.
30. Jamshidi Y, Snieder H, Ge D, Spector TD, O'Dell SD. The SH2B gene is associated with serum leptin and body fat in normal female twins. *Obesity (Silver Spring).* 2007 Jan;15(1):5-9.
31. Kaminsky ZA, Tang T, Wang SC, Ptak C, Oh GH, Wong AH, et al. DNA methylation profiles in monozygotic and dizygotic twins. *Nat Genet.* 2009 Feb;41(2):240-5.
32. Johannes F, Colot V, Jansen RC. Epigenome dynamics: a quantitative genetics perspective. *Nat Rev Genet.* 2008 Nov;9(11):883-90.
33. Symonds ME, Sebert SP, Hyatt MA, Budge H. Nutritional programming of the metabolic syndrome. *Nat Rev Endocrinol.* 2009 Nov;5(11):604-10.
34. Meaney MJ, Szyf M, Seckl JR. Epigenetic mechanisms of perinatal programming of hypothalamic-pituitary-adrenal function and health. *Trends Mol Med.* 2007 Jul;13(7):269-77.
35. Rumpersaud E, Mitchell BD, Naj AC, Pollin TI. Investigating parent of origin effects in studies of type 2 diabetes and obesity. *Curr Diabetes Rev.* 2008 Nov;4(4):329-39.
36. Kong A, Steinthorsdottir V, Masson G, Thorleifsson G, Sulem P, Besenbacher S, et al. Parental origin of sequence variants associated with complex diseases. *Nature.* 2009 Dec 17;462(7275):868-74.
37. Unoki H, Takahashi A, Kawaguchi T, Hara K, Horikoshi M, Andersen G, et al. SNPs in KCNQ1 are associated with susceptibility to type 2 diabetes in East Asian and European populations. *Nat Genet.* 2008 Sep;40(9):1098-102.
38. Yasuda K, Miyake K, Horikawa Y, Hara K, Osawa H, Furuta H, et al. Variants in KCNQ1 are associated with susceptibility to type 2 diabetes mellitus. *Nat Genet.* 2008 Sep;40(9):1092-7.
39. Li S, Zhao JH, Luan J, Luben RN, Rodwell SA, Khaw KT, et al. Cumulative effects and predictive value of common obesity-susceptibility variants identified by genome-wide association studies. *Am J Clin Nutr.* 2009 Oct 7.
40. Meigs JB, Shrader P, Sullivan LM, McAteer JB, Fox CS, Dupuis J, et al. Genotype score in addition to common risk factors for prediction of type 2 diabetes. *N Engl J Med.* 2008 Nov 20;359(21):2208-19.
41. Wray NR, Goddard ME, Visscher PM. Prediction of individual genetic risk of complex disease. *Curr Opin Genet Dev.* 2008 Jun;18(3):257-63.

42. Frazer KA, Murray SS, Schork NJ, Topol EJ. Human genetic variation and its contribution to complex traits. *Nat Rev Genet.* 2009 Apr;10(4):241-51.
43. Lindgren CM, Heid IM, Randall JC, Lamina C, Steinthorsdottir V, Qi L, et al. Genome-wide association scan meta-analysis identifies three Loci influencing adiposity and fat distribution. *PLoS Genet.* 2009 Jun;5(6):e1000508.
44. Dupuis J, Langenberg C, Prokopenko I, Saxena R, Soranzo N, Jackson AU, et al. New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. *Nat Genet.* Jan 17.
45. Herbert A, Gerry NP, McQueen MB, Heid IM, Pfeufer A, Illig T, et al. A common genetic variant is associated with adult and childhood obesity. *Science.* 2006 Apr 14;312(5771):279-83.
46. Dina C, Meyre D, Samson C, Tichet J, Marre M, Jouret B, et al. Comment on "A common genetic variant is associated with adult and childhood obesity". *Science.* 2007 Jan 12;315(5809):187; author reply
47. Loos RJ, Barroso I, O'Rahilly S, Wareham NJ. Comment on "A common genetic variant is associated with adult and childhood obesity". *Science.* 2007 Jan 12;315(5809):187; author reply
48. Rosskopf D, Bornhorst A, Rimbach C, Schwahn C, Kayser A, Kruger A, et al. Comment on "A common genetic variant is associated with adult and childhood obesity". *Science.* 2007 Jan 12;315(5809):187; author reply
49. Lyon HN, Emilsson V, Hinney A, Heid IM, Lasky-Su J, Zhu X, et al. The association of a SNP upstream of INSIG2 with body mass index is reproduced in several but not all cohorts. *PLoS Genet.* 2007 Apr 27;3(4):e61.
50. Andreasen CH, Mogensen MS, Borch-Johnsen K, Sandbaek A, Lauritzen T, Sorensen TI, et al. Non-replication of genome-wide based associations between common variants in INSIG2 and PFKP and obesity in studies of 18,014 Danes. *PLoS One.* 2008;3(8):e2872.
51. Boes E, Kollerits B, Heid IM, Hunt SC, Pichler M, Paulweber B, et al. INSIG2 polymorphism is neither associated with BMI nor with phenotypes of lipoprotein metabolism. *Obesity (Silver Spring).* 2008 Apr;16(4):827-33.
52. Chu X, Erdman R, Susek M, Gerst H, Derr K, Al-Agha M, et al. Association of morbid obesity with FTO and INSIG2 allelic variants. *Arch Surg.* 2008 Mar;143(3):235-40; discussion 41.
53. Franks PW, Jablonski KA, Delahanty LM, McAteer JB, Kahn SE, Knowler WC, et al. Assessing gene-treatment interactions at the FTO and INSIG2 loci on obesity-related traits in the Diabetes Prevention Program. *Diabetologia.* 2008 Dec;51(12):2214-23.
54. Hotta K, Nakamura M, Nakata Y, Matsuo T, Kamohara S, Kotani K, et al. INSIG2 gene rs7566605 polymorphism is associated with severe obesity in Japanese. *J Hum Genet.* 2008;53(9):857-62.
55. Orkunoglu-Suer FE, Gordish-Dressman H, Clarkson PM, Thompson PD, Angelopoulos TJ, Gordon PM, et al. INSIG2 gene polymorphism is associated with increased subcutaneous fat in women and poor response to resistance training in men. *BMC Med Genet.* 2008;9:117.
56. Reinehr T, Hinney A, Nguyen TT, Hebebrand J. Evidence of an influence of a polymorphism near the INSIG2 on weight loss during a lifestyle intervention in obese children and adolescents. *Diabetes.* 2008 Mar;57(3):623-6.
57. Wang HJ, Zhang H, Zhang SW, Pan YP, Ma J. Association of the common genetic variant upstream of INSIG2 gene with obesity related phenotypes in Chinese children and adolescents. *Biomed Environ Sci.* 2008 Dec;21(6):528-36.
58. Yang L, Wu Y, Li H, Yu Z, Li X, Liu Y, et al. Potential association of INSIG2 rs7566605 polymorphism with body weight in a Chinese subpopulation. *Eur J Hum Genet.* 2008 Jun;16(6):759-61.
59. Bressler J, Fornage M, Hanis CL, Kao WH, Lewis CE, McPherson R, et al. The INSIG2 rs7566605 genetic variant does not play a major role in obesity in a sample of 24,722 individuals from four cohorts. *BMC Med Genet.* 2009;10:56.
60. Ioannidis JP. Why most discovered true associations are inflated. *Epidemiology.* 2008 Sep;19(5):640-8.

61. Heid IM, Huth C, Loos RJ, Kronenberg F, Adamkova V, Anand SS, et al. Meta-analysis of the INSIG2 association with obesity including 74,345 individuals: does heterogeneity of estimates relate to study design? *PLoS Genet.* 2009 Oct;5(10):e1000694.
62. van Vliet-Ostaptchouk JV, Hofker MH, van der Schouw YT, Wijmenga C, Onland-Moret NC. Genetic variation in the hypothalamic pathways and its role on obesity. *Obes Rev.* 2009 Nov;10(6):593-609.
63. Saxena R, Hivert MF, Langenberg C, Tanaka T, Pankow JS, Vollenweider P, et al. Genetic variation in GIPR influences the glucose and insulin responses to an oral glucose challenge. *Nat Genet.* Jan 17.
64. Schadt EE. Molecular networks as sensors and drivers of common human diseases. *Nature.* 2009 Sep 10;461(7261):218-23.
65. Dimas AS, Deutsch S, Stranger BE, Montgomery SB, Borel C, Attar-Cohen H, et al. Common regulatory variation impacts gene expression in a cell type-dependent manner. *Science.* 2009 Sep 4;325(5945):1246-50.
66. Cheung VG, Spielman RS. Genetics of human gene expression: mapping DNA variants that influence gene expression. *Nat Rev Genet.* 2009 Sep;10(9):595-604.
67. Emilsson V, Thorleifsson G, Zhang B, Leonardson AS, Zink F, Zhu J, et al. Genetics of gene expression and its effect on disease. *Nature.* 2008 Mar 27;452(7186):423-8.
68. Leonardson AS, Zhu J, Chen Y, Wang K, Lamb JR, Reitman M, et al. The effect of food intake on gene expression in human peripheral blood. *Hum Mol Genet.* Jan 1;19(1):159-69.
69. Park PJ. ChIP-seq: advantages and challenges of a maturing technology. *Nat Rev Genet.* 2009 Oct;10(10):669-80.
70. Gstaiger M, Aebersold R. Applying mass spectrometry-based proteomics to genetics, genomics and network biology. *Nat Rev Genet.* 2009 Sep;10(9):617-27.
71. Stolk RP, Rosmalen JG, Postma DS, de Boer RA, Navis G, Slaets JP, et al. Universal risk factors for multifactorial diseases: LifeLines: a three-generation population-based study. *Eur J Epidemiol.* 2008;23(1):67-74.

